

Target Tailored Forest Damage Inventory (TFDI) - Damage by bark beetle in the county of Västernorrland and eastern Jämtland 2016

Dataset identifier: <https://doi.org/10.5878/thz3-ft06>

Contact: Cornelia Roberge, Department of Forest Resource Management, Swedish University of Agricultural Sciences

cornelia.roberge@slu.se

Report: Wulff, S. (2016). Nationell Riktad Skogsskadeinventering (NRS) 2016. Arbetsrapport 466. Institutionen för skoglig resurshushållning, Sveriges lantbruksuniversitet. <https://res.slu.se/id/publ/79768>

Instruction fieldwork: Separate file (Only in Swedish)

Data description:

Collected, assessed variables and calculated variables

PLOTDATA

Plotno

Date – Date of inventory (YYYYMMDD)

NFI year – Year when the NFI last time inventoried the sample plot before this inventory

Tract – Tract number

Sample plot – Sample plot number

Sub sample plot – Sub sample plot number (0=one plot)

County – 7 = Ångermanland; 8 = Medelpad; 5 = Jämtland

Inventory – Is the plot measured? 1=Yes; 2=No - sample plot not found; 3= No - Not forestland, a real change in land use class; 4=No - Whole plot clear cut; 5=No - Other reason (give explanation); 6=No – No Norway spruces left on the sample plot; 7=Yes – But only a few remaining trees; 8=Yes – But with a new stand edge (season 0 – 2) stand edge within 25 m

Plot area percent – Current proportion of sample plot

Plot area m² – Area of sample plot

Proportion of Norway spruce – Proportion of Norway spruce at plot level in tenths (NULL=Missing data)

Maturity class - Maturity class (see NFI instructions for description)

Edge - Distance from plot center to stand edge or gap in the forest (0=in the edge; 5=1-5 m; 10=6-10 m; 15=11-15 m; 20=16-20 m; 25=21-25 m; 26=>25 m; NULL=Missing data)

Edge time – Time when an stand edge within 25 m occurred (0=current season;
1=last year season; 2=season 2; 5=season 3-5; 10=older than 5 season; 99=no edge;
NULL=Missing data)

A season is from spring budding one year to the budding the next year.

Proportion of trees damaged 10m radius – Proportion of trees damaged by bark beetle at sample plot with 10m radius (NULL=Missing data)

Proportion of trees damaged 25m radius – Proportion of trees damaged by bark beetle at sample plot with 25m radius (NULL=Missing data)

Wind-felled trees season 0 – Number of wind thrown trees from season 0 (NULL=Missing data)

Wind-felled trees season 1 – Number of wind thrown trees from season 1 (NULL=Missing data)

Wind-felled trees fresh – Number of wind thrown trees with fresh inner bark (NULL=Missing data)

Wind-felled trees season older – Number of wind thrown trees older than season 1 (NULL=Missing data)

GPS-East Scrambled (SWEREF99, scrambled +/- 200 - 800 m)

GPS-North Scrambled (SWEREF99, scrambled +/- 200 - 800 m)

Other info – Other information free text in Swedish

TREEDATA

Plotno

Treeno

Within 10m – Within plot 10 m radius (1=yes; 0=no)

Dead – Dead tree (1=yes; 0=no)

Position – 1=standing tree; 2= Stump

Diameter – Diameter at breast height (mm)

Bark beetle – Damage by bark beetle species GBB=European spruce bark beetle (*Ips typographus*); DOB= Four eyed spruce bark beetle (*Polygraphus poligraphus*); GBBDOB=both European spruce bark beetle (*Ips typographus*) and Four eyed spruce bark beetle (*Polygraphus poligraphus*); ANN=Other bark beetle species;

Season – Damage season. A season is from spring budding one year to the budding the next year.
0=present season

Edge - Distance from tree to stand edge or gap in the forest (0=in the edge; 5=1-5 m; 10=6-10 m; 15=11-15 m; 20=16-20 m; 25=21-25 m; 26=>25 m) (NULL=Missing data)

Height – Estimated height (m) by height function RuneOlla (Ollas, R. 1980. Nya utbytesfunktioner för träd och bestånd. Forskningsstiftelsen Skogsarbeten, Ekonomi nr 5. In Swedish.)

Volume – Volume in m³ (Cubic meters standing volume)

Other info – Other information free text in Swedish

WIND-FELLED TREE DATA

Plotno

Treeno

Within 10m – Within plot 10 m radius (1=yes; 0=no)

Diameter – Diameter at breast height (mm)

Full length – Full length (1=yes; 0=no)

Length – If full length =1 than: Estimated height (m) by height function RuneOlla (Ollas, R. 1980. Nya utbytesfunktioner för träd och bestånd. Forskningsstiftelsen Skogsarbeten, Ekonomi nr 5. In Swedish.); If full length = 0 than measured length (m)

Top diameter – NULL if full length =1; If full length = 0 than measured diameter (mm)

Including stump – The wind-felled tree has a stump (1=yes; 0=no)

Infestation – The wind-felled tree infested with bark beetle (0 = no; 1=yes (only season 0))

Bark beetle – Damage by bark beetle species GBB=European spruce bark beetle (*Ips typographus*); DOB= Four eyed spruce bark beetle (*Polygraphus poligraphus*); ANN=Other bark beetle species; NULL=No attack of bark beetles

Deduction season – Wind-felled damage season. A season is from spring budding one year to the budding the next year. 0=present season; 1=last season; 2=two seasons ago; 3=two seasons ago or older

Volume – Volume in m³ (Cubic meters standing volume)

Other info – Other information free text in Swedish